

Sam--Formula

1. Into Blender Bottle, weigh out:
 - 480g of pre-mixed similac +
 - 16g of polycose +
 - 16g of similac powder
 - =16oz total
2. Each time of day is a different amount. Reference daily schedule sheet.
3. If you have to tube Sam instead of feed him orally, do the following:
 - if it is the 8:30am feed, do as you do at the 11:00 feed (Rate: 195, Dose: 100)
 - **measure out 4g of Boost/Pediasure + 126g of formula)

 - If it is the 1:30pm or 4:00pm feed do the following:
 - **measure out 3g of Boost/Pediasure + 100g of formula)

Sam fed:

- 8:30am (40g solids, 60g liquids)
- 11:00am (100g liquids)
- 1:30pm (40g solids, 60g liquids)
- 4:00pm (40g solids, 60g liquids)
- 6:15pm-- start overnight tube. Measure out 445cc total. Each night will be different; we are getting him used to more Pediasure.

- Thursday night:
 - 4oz boost/pediasure, rest formula. Program rate of 35cc/hr and dose 430cc. Measure 445 to make sure it doesn't run out.
- Friday night:
 - 5oz boost/pediasure, rest formula. Program rate of 35cc/hr and dose 430cc. Measure 445 to make sure it doesn't run out.
- Saturday night:
 - 6oz boost/pediasure, rest formula. Program rate of 35cc/hr and dose 430cc. Measure 445 to make sure it doesn't run out

Jack

Whole milk! Want him to drink ~2 cups each day. Give him milk before each meal, and during. I set him in high chair while I'm preparing/cutting up the food.

Behavior... If (when) he throws the cup, don't react. Let it sit on floor for ~3 minutes before picking up.

If he bangs the cup on table, take it away from him. Don't give back for ~3 minutes.